

Assessing the accuracy of the surface gravity determination in late-type stars with solar-like pulsators

Thierry Morel^{1*} and Andrea Miglio²

¹*Institut d'Astrophysique et de Géophysique, Université de Liège, Allée du 6 Août, Bât. B5c, 4000 Liège, Belgium*

²*School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT*

Accepted 2011 October 10. Received 2011 October 10; in original form 2011 September 7

ABSTRACT

The frequency of maximum oscillation power measured in dwarfs and giants exhibiting solar-like pulsations provides a precise, and potentially accurate, inference of the stellar surface gravity. An extensive comparison for about 40 well-studied pulsating stars with gravities derived using classical methods (ionization balance, pressure-sensitive spectral features or location with respect to evolutionary tracks) supports the validity of this technique and reveals an overall remarkable agreement with mean differences not exceeding 0.05 dex (although with a dispersion of up to ~ 0.2 dex). It is argued that interpolation in theoretical isochrones may be the most precise way of estimating the gravity by traditional means in nearby dwarfs. Attention is drawn to the usefulness of seismic targets as benchmarks in the context of large-scale surveys.

Key words: asteroseismology – stars: fundamental parameters – stars: late-type.

1 INTRODUCTION

It is notoriously difficult to accurately estimate the stellar surface gravity in late-type stars, with systematic differences of the order of 0.2 dex being commonplace depending on the technique used and its exact implementation. This large uncertainty surrounding $\log g$ limits the accuracy with which elemental abundances can be determined. This is especially the case for purely spectroscopic analyses where the determinations of the stellar parameters are intimately coupled. In such a case, the use of a model atmosphere with an inappropriate gravity adversely impacts on the estimation of the other parameters (i.e. effective temperature and microturbulence) and, ultimately, chemical abundances.

However, the properties of the p -mode pulsations exhibited by cool stars on the main sequence and during the red giant phase can be used to derive values that are precise to a level rivalling that obtained for eclipsing binaries. Although seismic gravities can also be derived using oscillation frequencies and frequency separations, here we only consider the frequency of maximum power, ν_{\max} , as a surface gravity indicator (see e.g. Kallinger et al. 2010a, for definition and further details on how this quantity can be derived). As first suggested by Brown et al. (1991) and recently discussed from a theoretical viewpoint by Belkacem et al. (2011), ν_{\max} is expected to scale as the acoustic cut-off frequency:

$$\frac{\nu_{\max}}{\nu_{\max, \odot}} = \left(\frac{M}{M_{\odot}} \right) \left(\frac{R}{R_{\odot}} \right)^{-2} \left(\frac{T_{\text{eff}}}{T_{\text{eff}, \odot}} \right)^{-1/2}. \quad (1)$$

This leads to

$$\log g = \log g_{\odot} + \log \left(\frac{\nu_{\max}}{\nu_{\max, \odot}} \right) + \frac{1}{2} \log \left(\frac{T_{\text{eff}}}{T_{\text{eff}, \odot}} \right). \quad (2)$$

This relation is largely insensitive to the effective temperature assumed ($\Delta T_{\text{eff}} = 100$ K leads to $\Delta \log g \sim 0.004$ dex only for Sun-like stars). On the other hand, ν_{\max} can usually be measured with an error below 5 per cent from high-quality time series (e.g. Kallinger et al. 2010a; Mosser et al. 2010). It follows that $\log g$ determined via equation (2) can be precise to better than 0.03 dex. If confirmed in terms of accuracy, this would be far better than what can be achieved by other means in single stars (except in stars with transiting planets, although this method heavily relies on evolutionary models; e.g. Torres, Winn & Holman 2008). Indeed, seismic gravities are beginning to be adopted in spectroscopic analyses as an alternative to values derived from traditional methods in order to narrow down the uncertainties in the other fundamental stellar parameters and chemical abundances (e.g. Batalha et al. 2011).

The high accuracy of the gravities obtained from asteroseismology is supported by a comparison with values estimated using completely independent techniques [e.g. as shown in the case of a few binaries by Bruntt et al. (2010), as well as in red giant cluster members by Stello et al. (2011) and Miglio et al. (2011)]. However, the validity of the scalings relating the stellar parameters (mass, radius) and the seismic observables has yet to be thoroughly investigated for stars occupying different parts of the Hertzsprung-Russell diagram and having various properties in terms of metallicity and activity level, for instance. This work is an effort towards this goal (see also Miglio 2011) and also aims at drawing attention to the usefulness of seismic targets for validation purposes in the context of large-scale stellar surveys.

*E-mail: morel@astro.ulg.ac.be

2 A SAMPLE OF WELL-STUDIED STARS WITH A PRECISE SEISMIC GRAVITY

About 40 bright, well-studied solar-like and red giant stars have an accurate estimate of the frequency of maximum power, either from ground-based radial-velocity monitoring or from ultraprecise photometric observations from space (Table 1). These have been used, along with mean literature T_{eff} values (see below) and assuming $\nu_{\text{max}, \odot} = 3100 \mu\text{Hz}$, to compute the seismic gravities (the exact choice of $\nu_{\text{max}, \odot}$, which has an uncertainty of $\sim 50 \mu\text{Hz}$, has a negligible impact on our results). The temperatures adopted are marginally higher than those derived from angular diameter and bolometric flux measurements (Bruntt et al. 2010): $(\Delta T_{\text{eff}}) = +44 \pm 56 \text{ K}$ (1σ , 10 stars). Adopting these values would lead to negligible differences in the seismic $\log g$ (well below 0.01 dex). The uncertainty in ν_{max} , which is the main source of error, is often not quoted in the original literature source or its estimation relies on widely different criteria and assumptions. It is therefore impossible to properly account for the star-to-star differences in the data quality and provide a homogeneous set of uncertainties. Adopting various procedures for the determination of ν_{max} and taking into account the different signal realizations arising from the stochastic nature of the oscillations, Hekker et al. (2011) inferred an uncertainty in the range 1–10 per cent for stars observed by the *Kepler* mission. Based on the type of data collected for the stars in our sample, we estimate a typical uncertainty of 5 per cent. This translates into an error in the seismic gravities of ~ 0.03 dex only. These figures are supported by a comparison with the values for the three stars in binaries with dynamical masses and interferometric radii (Procyon A and α Cen A+B; Bruntt et al. 2010): the gravities agree to within 0.02 dex.

3 THE CLASSICAL GRAVITY DIAGNOSTICS USED IN COOL STARS PUT TO THE TEST

As the stars in Table 1 are amongst the brightest in the sky and are even sometimes regarded as standards (e.g. α Boo or Procyon A), a large number of independent determinations from classical techniques can be found in the literature. This offers an opportunity to empirically assess the reliability of the most popular gravity diagnostics used in cool stars: ionization balance of a given chemical species (usually iron), fitting the wings of pressure-sensitive, neutral metal lines or interpolation in theoretical isochrones. Discrepancies between the various indicators are known to exist in the case of unresolved, single-lined binaries (Fuhrmann et al. 2011), but this should not be a concern in our sample. These three approaches suffer to different extents from drawbacks. First, the values obtained from ionization equilibrium are strongly dependent on the atmospheric structure adopted (granulation is ignored in 1D models) and can be biased by non-local thermodynamic equilibrium (non-LTE) effects, which become generally more important in stars with extended atmospheres and/or metal poor [see e.g. the calculations of Mashonkina et al. (2011) applied to some stars in our sample]. On the other hand, values obtained from fitting the wings of pressure-sensitive lines are generally affected by quite large uncertainties related, for instance, to blends and difficulties in continuum placement (e.g. Bruntt et al. 2010). Finally, although for very nearby stars parallaxes and reddening are not a major concern, interpolation in theoretical isochrones is strongly model-dependent and may suffer from degeneracy problems (as a result, the applicability of this method for stars on the red giant branch is limited).


The T_{eff} , [Fe/H] and $\log g$ literature values for the stars in Table 1 were primarily extracted from the PASTEL catalogue

Table 1. Values of the frequency of maximum power, ν_{max} , for the stars in our sample. The typical uncertainty is 5 per cent (see text). The original references for the seismic data are given. When not explicitly quoted in these papers, the ν_{max} values were taken from Bruntt et al. (2010), Kallinger et al. (2010a) or Mosser et al. (2010). The value for 18 Sco was computed from the original data.

Name	ν_{max} (μHz)	Reference	Name	ν_{max} (μHz)	Reference		
Dwarfs							
HD 2151	β Hyi	1000	Kjeldsen et al. (2008)	HD 165341	70 Oph A	4500	Carrier & Eggenberger (2006)
HD 10700	τ Cet	4490	Teixeira et al. (2009)	HD 170987		930	Mathur et al. (2010)
HD 17051	ι Hor	2700	Vauclair et al. (2008)	HD 175726		2000	Mosser et al. (2009)
HD 20010	α For	1100	Kjeldsen et al. (2008)	HD 181420		1500	Barban et al. (2009)
HD 23249	δ Eri	700	Bouchy & Carrier (2003)	HD 181906		1912	García et al. (2009)
HD 49385		1013	Deheuvels et al. (2010)	HD 190248	δ Pav	2300	Kjeldsen et al. (2008)
HD 49933		1657	Kallinger et al. (2010b)	HD 203608	γ Pav	2600	Mosser et al. (2008)
HD 52265		2090	Ballot et al. (2011)	HD 210302	τ Psa	1950	Bruntt et al. (2010)
HD 61421	Procyon A	1000	Arentoft et al. (2008)	Subgiants and giants			
HD 63077	171 Pup	2050	Bruntt et al. (2010)	HD 71878	β Vol	51	Stello et al. (2009)
HD 102870	β Vir	1400	Carrier et al. (2005a)	HD 100407	ξ Hya	92.3	Frandsen et al. (2002)
HD 121370	η Boo	750	Carrier, Eggenberger & Bouchy (2005b)	HD 124897	α Boo	3.47	Tarrant et al. (2007)
HD 128620	α Cen A	2400	Kjeldsen et al. (2008)	HD 146791	ϵ Oph	53.5	Kallinger et al. (2008)
HD 128621	α Cen B	4100	Kjeldsen et al. (2008)	HD 153210	κ Oph	35	Stello et al. (2009)
HD 139211	HR 5803	2800	Carrier, Eggenberger & Leyder (2008)	HD 161096	β Oph	46	Kallinger et al. (2010a)
HD 142860	γ Ser	1600	Kjeldsen et al. (2008)	HD 163588	ξ Dra	36	Stello et al. (2009)
HD 146233	18 Sco	3170	Bazot et al. (2011)	HD 168723	η Ser	125	Barban et al. (2004)
HD 150680	ζ Her A	700	Martić et al. (2001)	HD 188512	β Aql	410	Kjeldsen et al. (2008)
HD 160691	μ Ara	2000	Bouchy et al. (2005)	HD 211998	ν Ind	313	Bedding et al. (2006)
HD 161797	μ Her	1200	Bonanno et al. (2008)	M67 S1305		208.9	Kallinger et al. (2010a)

Table 2. Mean effective temperature, iron content and surface gravities from the four different methods for the stars in Table 1. The complete table is available online (see Supporting Information).

Name	T_{eff} (K)	[Fe/H]	Seismology	log g			
				Ionization	Wings	Isochrone	
Dwarfs							
HD 2151	β Hyi	5829 ± 107 (10)	-0.09 ± 0.12 (11)	3.95	4.02 ± 0.18 (6)	3.76 ± 0.15 (1)	3.98 ± 0.10 (8)
...

**Figure 1.** Difference between the seismic log g values and those obtained through ionization balance of iron (*top panels*), fitting of the wings of pressure-sensitive lines (*middle panels*) and isochrone fitting (*bottom panels*), as a function of the seismic gravities, effective temperature and metallicity. Representative error bars are indicated.

(Soubiran et al. 2010), but were supplemented by data from several missing sources after a careful inspection of the vast literature for these objects. Only studies published after 1990 were considered, as older ones may be based on poor-quality data or inadequate model atmospheres. Each original reference was inspected to evaluate the method used for the log g determination. In some instances, a single value was quoted in PASTEL, whereas estimates based on different techniques were reported in the original paper [e.g. Santos et al. (2005) where the gravities estimated from isochrone fitting are missing]. These values were added. Finally, duplicate entries from the same authors were omitted; only the value in the most recent paper was used. This roughly totals to 360 individual measurements from 80 independent literature sources. The results are presented in Table 2.

The comparison between the seismic log g values and those obtained through traditional techniques is shown in Fig. 1. Overall, there is a remarkably good agreement with systematic differences not exceeding 0.04 dex on average. The significant 1σ dispersion of up to 0.19 dex with respect to the reference seismic value may have been expected considering the heterogeneous nature of the data and the diversity of analyses performed. It remains to be seen, however, if part of the observed scatter is not due to an intrinsic dispersion of equation (2), which is based on a simple scaling of v_{max} with the acoustic cut-off frequency in the atmosphere. By averaging results from a large number of independent studies (as is the case here for the ionization and isochrone gravities, but *not* for the strong-line ones), one can hope that the systematic errors partly cancel out and that the mean offset with respect to the seismic values provides a better appraisal of the true accuracy of the method. It should be kept in mind that the systematic differences which may exist between the

various analyses might not be completely related to the technique used, but instead to other assumptions in the modelling (especially the T_{eff} scale). The results of the studies with the highest number of measurements are shown in Fig. 2. As can be seen, several of them can be in error by more than a factor of 2.

It can readily be seen in Fig. 1 that the scatter is lower for the gravities estimated from isochrone fitting. The same conclusion holds when considering for each star the average of the measurements obtained using a given method (Fig. 3),¹ especially when one excludes the evolved objects ($\log g < 3.2$) for which the determination through the position of the star with respect to evolutionary tracks is ill defined. In that case, the difference scatter is a mere ~ 15 per cent: $\langle \Delta \log g \rangle = -0.006 \pm 0.065$ dex (1σ , 29 stars). The evolutionary tracks used mostly fall in two categories differing in their physical ingredients (e.g. treatment of convection): either those from the Geneva (e.g. Schaller et al. 1992) or from the Padova (e.g. Bertelli et al. 1994) group. None the less, the gravity determination seems fairly robust against the choice of the set of models adopted (see Allende Prieto et al. 1999). Although this method is generally the most precise, it must be stressed that the mean difference with respect to the seismic gravities is less than 0.05 dex irrespective of the technique used.

No clear trends are discernible as a function of log g , T_{eff} or [Fe/H]. An underestimation of log g through ionization balance may be expected for very metal-poor stars ([Fe/H] $\lesssim -1$) because of non-LTE effects (Allende Prieto et al. 1999). We only have one such star

¹ An unweighted mean has been used because of the difficulty in assessing the quality of the studies performed and the inhomogeneous nature of the quoted uncertainties (which are often only internal and, as a result, appear unrealistically small).




Figure 2. As Fig. 1, but for the key studies only. Black, open circles: Allende Prieto & Lambert (1999) and Allende Prieto et al. (2004); cyan, filled circles: Bruntt (2009) and Bruntt et al. (2010); yellow crosses: Gonzalez & Wallerstein (1998), Gonzalez et al. (2001) and Gonzalez, Carlson & Tobin (2010); red, filled triangles: Luck & Heiter (2006, 2007); magenta, open triangles: Ramírez, Allende Prieto & Lambert (2007); blue, filled squares: Santos, Israelian & Mayor (2001, 2004) and Santos et al. (2005); green, open squares: Takeda et al. (2005, 2007) and Takeda, Sato & Murata (2008).




Figure 3. As Fig. 1, but with the data averaged on a star-to-star basis.

in our sample (ν Ind), but the ionization gravity does not appear discrepant. The $\log g$ values are systematically underestimated in the dwarfs by up to 0.3 dex when fitting the wings of strong metal lines. However, the bulk of the data comes from a single source (Bruntt et al. 2010) and large line-to-line differences are observed (a weighted mean has been used here).

4 CONCLUSIONS AND PERSPECTIVES

The good agreement between the gravities inferred from asteroseismology and from classical methods supports the applicability of the scaling law linking $\log g$ and ν_{\max} for stars spanning a relatively wide range in temperature and evolutionary status, although its validity in the low-metallicity and low-gravity regimes cannot be meaningfully investigated here owing to the limited number of objects. For stars within the parameter space investigated here, our study hence supports the use of the seismic gravities as input in

spectroscopic analyses (e.g. Batalha et al. 2011). Gravities relying on asteroseismic information would be especially valuable for red giants in view of the fundamental difficulties plaguing the classical techniques in that case (significant departures from LTE, lack of sensitivity of strong metal lines and degeneracy of isochrone fitting). The spectroscopic gravities of the faint red giant *CoRoT* targets are indeed found to be affected by large errors (Morel, Miglio & Valentini 2011). Seismic information is now available for hundreds of stars in the *CoRoT* and *Kepler* fields (see e.g. Verner et al. 2011). It is therefore reasonable to assume that similar consistency checks as those presented in this Letter will be extended to much larger samples in the future.

A comparison with data for eclipsing binaries has already illustrated the power of isochrone fitting as gravity indicator (Allende Prieto & Lambert 1999), and our study indeed identifies it as being the most precise classical method for nearby dwarfs. The release of the *Gaia* parallaxes will offer the opportunity to apply this

technique to much fainter magnitudes, although reddening will remain an issue.

Several large-scale spectroscopic surveys are presently conducted or are being planned (e.g. Radial Velocity Experiment, Apache Point Observatory Galactic Evolution Experiment or the follow-up of the *Gaia* mission). The pipelines developed for that purpose should be able to recover the parameters determined for a set of training stars through completely different and, as much as possible, model-independent methods before embarking on the automatic analysis of large samples of potentially faint objects. The seismic gravities can hence constitute a valuable piece of information in this context. Of particular interest in this respect are the stars with an accurate T_{eff} and $\log g$ estimate from interferometric and seismic observations, respectively (see Bruntt et al. 2010).

ACKNOWLEDGMENTS

TM acknowledges financial support from Belspo for contract PRODEX GAIA-DPAC and is indebted to S. Vauclair for drawing his attention to the accuracy of the seismic gravities. We would like to thank the anonymous referee for valuable comments, A.-M. Broomhall for recomputing the ν_{max} value of 18 Sco and M. Valentini for useful discussions.

REFERENCES

- Allende Prieto C., Lambert D. L., 1999, *A&A*, 352, 555
 Allende Prieto C., García López R. J., Lambert D. L., Gustafsson B., 1999, *ApJ*, 527, 879
 Allende Prieto C., Barklem P. S., Lambert D. L., Cunha K., 2004, *A&A*, 420, 183
 Arentoft T. et al., 2008, *ApJ*, 687, 1180
 Ballot J. et al., 2011, *A&A*, 530, A97
 Barban C. et al., 2004, in Wilson A., ed, *Proc. Soho 14/Gong 2004 Workshop, Helio- and Asteroseismology; towards a golden future*, ESA SP-559. ESA, Noordwijk, p. 113
 Barban C. et al., 2009, *A&A*, 506, 51
 Batalha N. M. et al., 2011, *ApJ*, 729, 27
 Bazot M. et al., 2011, *A&A*, 526, L4
 Bedding T. R. et al., 2006, *ApJ*, 647, 558
 Belkacem K., Goupil M. J., Dupret M. A., Samadi R., Baudin F., Noels A., Mosser B., 2011, *A&A*, 530, A142
 Bertelli G., Bressan A., Chiosi C., Fagotto F., Nasi E., 1994, *A&AS*, 106, 275
 Bonanno A., Benatti S., Claudi R., Desidera S., Gratton R., Leccia S., Paternò L., 2008, *ApJ*, 676, 1248
 Bouchy F., Carrier F., 2003, *Ap&SS*, 284, 21
 Bouchy F., Bazot M., Santos N. C., Vauclair S., Sosnowska D., 2005, *A&A*, 440, 609
 Brown T. M., Gilliland R. L., Noyes R. W., Ramsey L. W., 1991, *ApJ*, 368, 599
 Bruntt H., 2009, *A&A*, 506, 235
 Bruntt H. et al., 2010, *MNRAS*, 405, 1907
 Carrier F., Eggenberger P., 2006, *A&A*, 450, 695
 Carrier F., Eggenberger P., D'Alessandro A., Weber L., 2005a, *New Astron.*, 10, 315
 Carrier F., Eggenberger P., Bouchy F., 2005b, *A&A*, 434, 1085
 Carrier F., Eggenberger P., Leyder J.-C., 2008, *J. Phys. Conf. Ser.*, 118, 012047
 Deheuvels S. et al., 2010, *A&A*, 515, A87
 Frandsen S. et al., 2002, *A&A*, 394, L5
 Fuhrmann K., Chini R., Hoffmeister V. H., Stahl O., 2011, *MNRAS*, 415, 1240
 García R. A. et al., 2009, *A&A*, 506, 41
 Gonzalez G., Wallerstein G., 1998, *AJ*, 116, 765
 Gonzalez G., Laws C., Tyagi S., Reddy B. E., 2001, *AJ*, 121, 432
 Gonzalez G., Carlson M. K., Tobin R. W., 2010, *MNRAS*, 403, 1368
 Hekker S. et al., 2011, *A&A*, 525, A131
 Kallinger T. et al., 2008, *A&A*, 478, 497
 Kallinger T. et al., 2010a, *A&A*, 509, A77
 Kallinger T., Gruberbauer M., Guenther D. B., Fossati L., Weiss W. W., 2010b, *A&A*, 510, A106
 Kjeldsen H. et al., 2008, *ApJ*, 682, 1370
 Luck R. E., Heiter U., 2006, *AJ*, 131, 3069
 Luck R. E., Heiter U., 2007, *AJ*, 133, 2464
 Martić M., Lebrun J. C., Schmitt J., Appourchaux T., Bertaux J. L., 2001, in Wilson A., ed, *Proc. Soho 10/Gong 2000 Workshop, Helio- and Asteroseismology: at the dawn of the millenium*, ESA SP-464. ESA, Noordwijk, p. 431
 Mashonkina L., Gehren T., Shi J.-R., Korn A. J., Grupp F., 2011, *A&A*, 528, A87
 Mathur S. et al., 2010, *A&A*, 518, A53
 Miglio A., 2011, preprint (arXiv:1108.4555)
 Miglio A. et al., 2011, *MNRAS*, preprint (arXiv:1109.4376)
 Morel T., Miglio A., Valentini M., 2011, *J. Phys. Conf. Ser.*, in press
 Mosser B., Deheuvels S., Michel E., Thévenin F., Dupret M. A., Samadi R., Barban C., Goupil M. J., 2008, *A&A*, 488, 635
 Mosser B. et al., 2009, *A&A*, 506, 33
 Mosser B. et al., 2010, *A&A*, 517, A22
 Ramírez I., Allende Prieto C., Lambert D. L., 2007, *A&A*, 465, 271
 Santos N. C., Israelian G., Mayor M., 2001, *A&A*, 373, 1019
 Santos N. C., Israelian G., Mayor M., 2004, *A&A*, 415, 1153
 Santos N. C., Israelian G., Mayor M., Bento J. P., Almeida P. C., Sousa S. G., Ecuivillon A., 2005, *A&A*, 437, 1127
 Schaller G., Schaerer D., Meynet G., Maeder A., 1992, *A&AS*, 96, 269
 Soubiran C., Le Campion J.-F., Cayrel de Strobel G., Caillo A., 2010, *A&A*, 515, A111
 Stello D., Chaplin W. J., Basu S., Elsworth Y., Bedding T. R., 2009, *MNRAS*, 400, L80
 Stello D. et al., 2011, *ApJ*, 739, 13
 Takeda Y., Ohkubo M., Sato B., Kambe E., Sadakane K., 2005, *PASJ*, 57, 27
 Takeda Y., Kawanamoto S., Honda S., Ando H., Sakurai T., 2007, *A&A*, 468, 663
 Takeda Y., Sato B., Murata D., 2008, *PASJ*, 60, 781
 Tarrant N. J., Chaplin W. J., Elsworth Y., Spreckley S. A., Stevens I. R., 2007, *MNRAS*, 382, L48
 Teixeira T. C. et al., 2009, *A&A*, 494, 237
 Torres G., Winn J. N., Holman M. J., 2008, *ApJ*, 677, 1324
 Vauclair S., Laymand M., Bouchy F., Vauclair G., Hui Bon Hoa A., Charpinet S., Bazot M., 2008, *A&A*, 482, L5
 Verner G. A. et al., 2011, *ApJ*, 738, L28

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

Table S1. Mean effective temperature, iron content and surface gravities from the four different methods for the stars in Table 1. The error bars are the quadratic sum of the standard deviation of the individual measurements and the typical uncertainty in the parameter determination (80 K for T_{eff} , 0.1 dex for [Fe/H], 0.1 dex for the ionization and isochrone gravities, and 0.15 dex for the strong-line gravities). The numbers in brackets are the number of measurements. The typical uncertainty in the seismic $\log g$ is 0.03 dex (see text).

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.